Focus on Rectal Mass

Seon-Young Park, Hyun-Soo Kim

Division of Gastroenterology, Department of Internal Medicine, Chonnam National University Medical School, Gwangju, Korea

Received May 14, 2021, Revised May 26, 2021, Accepted May 26, 2021

Corresponding author: Seon-Young Park, E-mail: drpsy@naver.com, <https://orcid.org/0000-0002-0962-5977>

QUESTION: A 47-year-old man was admitted to a hospital for evaluation of rectal mass. He had been diagnosed with ulcerative colitis (UC) 10 years ago (Fig. 1), treated with intravenous steroid and remained in remission with oral mesalazine. He had a history of endoscopic mucosal

resection (EMR) for rectal polyp of 2 cm in diameter 5 years ago, which pathologic finding suggested hyperplastic polyp with inflammatory change (Fig. 2). Follow-up colonoscopy showed an about 3 cm sized recurrent rectal mass around the previous EMR site (Fig. 3). What is the diagnosis of rectal

Fig. 1. Colonoscopic findings at the diagnosis of ulcerative colitis.

Fig. 2. Rectal polyp. (A) Rectal polyp at the symptomatic remission with intravenous steroid. (B, C) Follow-up colonoscopy showed a rectal polyp around fibrotic mucosa, which was removed by endoscopic mucosal resection.

Fig. 3. Follow-up colonoscopy 10 years after initial diagnosis.

mass?

ANSWER: Rectal cancer in patients with long-standing ulcerative colitis.

REVIEW: Long-standing UC is associated with an increased risk of dysplasia or cancer, which develop through a different pathways compared to those in sporadic cancer [1]. In this patient with UC of duration longer than 10 years, rectal cancer (pT3N2Mx) was diagnosed. The pathomechanism is secondary to long-standing inflammation and associated with the duration and extent of diseases. Current guidelines describe starting surveillance colonoscopy 6 to 10 years after the diagnosis of UC [1-3]. The rate of missed malignancy in patients with UC is not insignificant and underestimates the importance of an effective surveillance program [4], which depends on many factors; patient compliance, adequate bowel preparation, adequate mucosal sampling or use of advanced endoscopic imaging for appropriate recognition of abnormal lesions. Debates still exists regarding the optimal strategy for determining which patients with early dysplasia can be managed endoscopically and which require radical surgery. While the optimal management of low grade dysplasia continues to be debated, the finding of high

grade dysplasia or colorectal cancer usually warrants surgical resection [5].

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

AUTHOR'S CONTRIBUTIONS

Conceptualization: Hyun-Soo Kim. Writing draft and supervision: Seon-Young Park.

ORCID

Seon-Young Park, <https://orcid.org/0000-0002-0962-5977>

Hyun-Soo Kim, <https://orcid.org/0000-0003-4834-0496>

REFERENCES

1. Gaidos JK, Bickston SJ. How to optimize colon cancer surveillance in inflammatory bowel disease patients. *Inflamm Bowel Dis* 2016;22:1219-1230.
2. Lamb CA, Kennedy NA, Raine T, et al. British Society of Gastroenterology consensus guidelines on the management of inflammatory bowel disease in adults. *Gut* 2019;68(Suppl 3):s1-s106.
3. Shergill AK, Lightdale JR, Bruining DH, et al.; American Society for Gastrointestinal Endoscopy Standards of Practice Committee. The role of endoscopy in inflammatory bowel disease. *Gastrointest Endosc* 2015;81:1101-1121.e1-e13.
4. Wang YR, Cangemi JR, Loftus EV Jr, Picco MF. Rate of early/missed colorectal cancers after colonoscopy in older patients with or without inflammatory bowel disease in the United States. *Am J Gastroenterol* 2013;108:444-449.
5. Huang LC, Merchea A. Dysplasia and cancer in inflammatory bowel disease. *Surg Clin North Am* 2017;97:627-639.